Technical advice

Neuguinea Impatiens IMPACIO®

Describtion:
When breeding pac Impacio® ourselves, we select for uniformity, good branching and high robustness against environmental factors. Impacio® offers a wide range of colours with many large single flowers.

Potting:
Week - 12
Pot 10,5 – 13 cm

Sales:
- Starts end of april

Substrate:
- Potting-Substrate
- pH 5,5 – 6,5

Fertilizer:
Low nutrient need: Do no use depot fertilizer. EC max: 1,2 to 1,8. High salt content leads to leaf damages.

Temperature:
Moderate temperature need. directly after potting daily average temperature 18°C, thereafter with DAT at 14°C. Hardening at 10°C.

Light:
After roots have established you can give full light during day.

Flowers:
End of april

Growth regulators:
Bonzi 2 ppm

Pinching:
Soft pinch, 6 -7 days after potting.
Further pinch may be necessary depending on growth.

Diseases / pests:
Botrytis, pythiam, rizoctionia, aphids, virus.

Spacing:
20 - 25 plants / sqm.

Elsner pac® Vertriebsgesellschaft mbH | Am Fiebig 14 | 01561 Thiendorf
Phone: +49 35248 3991-0 | E-Mail: info@pac-elsner.com

PASSION
AND COMPETENCE