Kulturempfehlung

Ostespermum

Describtion:

Our Osteospermums represent a range of varieties that grow well and branch profusely, producing large, elegant flowers. It is uniform in growth and flowering time and requires little growth inhibitor.

Potting:

Week 2 - 14
Pot 10,5 – 13 cm
Cold cultivation over winter possible.
Potting from week 45 onwards

Sales:

- Early march

Substrate:

- Potting-Substrate
- pH 5,5 – 6,5

Fertilizer:

High nutrient need: with continuous fertigation 0,08 % to 0,1 % or weekly fertigation 1-2 times per week 0,2 % to 0,3 %

Temperature:

Moderate temperature need. After potting start with daily average temperature 18°C, thereafter DAT 16°C.
Cold crop cultivation and overwintering is highly recommended.
Cool Morning and Diff/Drop Strategy recommended.

Light:

Full light can be given after root establishment

Flowers:

Starts in march

Growth regulators:

Bonzi 6 – 8 ppm, Alar 2500 – 4000 ppm

Pinching:

Soft pinch one week after potting. Further pinch may be necessary .

Krankheiten/Schädlinge:

Thrips, white fly, leaf miner, powdery mildew.

Flächenbelegung:

45-60 plants / sqm

Elsner pac® Vertriebsgesellschaft mbH | Am Fiebig 14 | 01561 Thiendorf
Phone: +49 35248 3991-0 | E-Mail: info@pac-elsner.com

PASSION
AND COMPETENCE